

## SMARANDACHE HYPER ALGEBRAS

YOUNG BAE JUN AND EUN HWAN ROH\*

Received July 12, 2005

ABSTRACT. The notion of Smarandache hyper  $I$ -algebra and Smarandache hyper  $K$ -algebra are introduced, and related properties are investigated.

## 1. INTRODUCTION

Generally, in any human field, a *Smarandache Structure* on a set  $A$  means a weak structure  $\mathbf{W}$  on  $A$  such that there exists a proper subset  $B$  of  $A$  which is embedded with a strong structure  $\mathbf{S}$ . In [5], W. B. Vasantha Kandasamy studied the concept of Smarandache groupoids, subgroupoids, ideal of groupoids, semi-normal subgroupoids, Smarandache Bol groupoids and strong Bol groupoids and obtained many interesting results about them. Smarandache semigroups are very important for the study of congruences, and it was studied by R. Padilla [4]. In this paper, we introduce the notion of Smarandache hyper  $K$ - and  $I$ -algebras, and investigate its properties.

## 2. PRELIMINARIES

We include some elementary aspects of hyper  $K$ -algebras that are necessary for this paper, and for more details we refer to [1] and [6]. Let  $H$  be a non-empty set endowed with a hyper operation “ $\circ$ ”, that is,  $\circ$  is a function from  $H \times H$  to  $\mathcal{P}^*(H) = \mathcal{P}(H) \setminus \{\emptyset\}$ . For two subsets  $A$  and  $B$  of  $H$ , denote by  $A \circ B$  the set  $\bigcup_{a \in A, b \in B} a \circ b$ .

By a *hyper BCK-algebra* we mean a non-empty set  $H$  endowed with a hyperoperation “ $\circ$ ” and a constant  $0$  satisfying the following axioms:

- (HK1)  $(x \circ z) \circ (y \circ z) \ll x \circ y$ ,
- (HK2)  $(x \circ y) \circ z = (x \circ z) \circ y$ ,
- (HK3)  $x \circ H \ll \{x\}$ ,
- (HK4)  $x \ll y$  and  $y \ll x$  imply  $x = y$ ,

for all  $x, y, z \in H$ , where  $x \ll y$  is defined by  $0 \in x \circ y$  and for every  $A, B \subseteq H$ ,  $A \ll B$  is defined by  $\forall a \in A, \exists b \in B$  such that  $a \ll b$ .

By a *hyper I-algebra* we mean a non-empty set  $H$  endowed with a hyper operation “ $\circ$ ” and a constant  $0$  satisfying the following axioms:

- (H1)  $(x \circ z) \circ (y \circ z) < x \circ y$ ,
- (H2)  $(x \circ y) \circ z = (x \circ z) \circ y$ ,
- (H3)  $x < x$ ,
- (H4)  $x < y$  and  $y < x$  imply  $x = y$

---

2000 *Mathematics Subject Classification.* 06F35, 03G25.

*Key words and phrases.* Smarandache hyper  $I$ -algebra, Smarandache hyper  $K$ -algebra,  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal,  $\Omega$ -Smarandache hyper  $(<, \in)$ -ideal,  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal,  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \sqsubseteq)$ -ideal.

\*Corresponding author: Tel. +82 55 740 1232.

for all  $x, y, z \in H$ , where  $x < y$  is defined by  $0 \in x \circ y$  and for every  $A, B \subseteq H$ ,  $A < B$  is defined by  $\exists a \in A$  and  $\exists b \in B$  such that  $a < b$ . If a hyper  $I$ -algebra  $(H, \circ, 0)$  satisfies an additional condition:

(H5)  $0 < x$  for all  $x \in H$ ,

then  $(H, \circ, 0)$  is called a *hyper  $K$ -algebra* (see [1]).

Note that every hyper  $BCK$ -algebra is a hyper  $K$ -algebra.

In a hyper  $I$ -algebra  $H$ , the following hold (see [1, Proposition 3.4]):

- (a1)  $(A \circ B) \circ C = (A \circ C) \circ B$ ,
- (a2)  $A \circ B < C \Leftrightarrow A \circ C < B$ ,
- (a3)  $A \subseteq B$  implies  $A < B$

for all nonempty subsets  $A, B$  and  $C$  of  $H$ .

In a hyper  $K$ -algebra  $H$ , the following hold (see [1, Proposition 3.6]):

- (a4)  $x \in x \circ 0$  for all  $x \in H$ .

### 3. SMARANDACHE HYPER ALGEBRAS AND SMARANDACHE HYPER IDEALS

**Definition 3.1.** A *Smarandache hyper  $I$ -algebra* is defined to be a hyper  $I$ -algebra  $(H, \circ, 0)$  in which there exists a proper subset  $\Omega$  of  $H$  such that  $(\Omega, \circ, 0)$  is a non-trivial hyper  $K$ -algebra.

**Example 3.2.** Let  $H = \{0, 1, 2\}$  and define an hyper operations “ $\circ_1$ ” and “ $\circ_2$ ” on  $H$  as follows:

$\circ_1$	0	1	2
0	$\{0\}$	$\{0\}$	$\{2\}$
1	$\{1\}$	$\{0\}$	$\{2\}$
2	$\{2\}$	$\{2\}$	$\{0, 2\}$

Table a1

$\circ_2$	0	1	2
0	$\{0\}$	$\{0, 1\}$	$\{2\}$
1	$\{1\}$	$\{0\}$	$\{2\}$
2	$\{2\}$	$\{0\}$	$\{0, 1, 2\}$

Table a2

Then  $(H, \circ_1, 0)$  and  $(H, \circ_2, 0)$  are Smarandache hyper  $I$ -algebras.

**Example 3.3.** Let  $H = \{0, a, b\}$  and define an hyper operation “ $\circ$ ” on  $H$  by the following Cayley table:

$\circ$	0	a	b
0	$\{0, a, b\}$	$\{a, b\}$	$\{a, b\}$
a	$\{0, a, b\}$	$\{0, a, b\}$	$\{a, b\}$
b	$\{0, a, b\}$	$\{a, b\}$	$\{0, a, b\}$

Table a4

Then  $(H, \circ, 0)$  is not a Smarandache hyper  $I$ -algebra since  $(\Omega_1 = \{0, a\}, \circ, 0)$  and  $(\Omega_2 = \{0, b\}, \circ, 0)$  are not hyper  $K$ -algebras.

**Definition 3.4.** A *Smarandache hyper  $K$ -algebra* is defined to be a hyper  $K$ -algebra  $(H, \circ, 0)$  in which there exists a proper subset  $\Omega$  of  $H$  such that  $(\Omega, \circ, 0)$  is a non-trivial hyper  $BCK$ -algebra.

**Example 3.5.** Let  $H = \{0, a, b, c\}$  and define an hyper operation “ $\circ$ ” on  $H$  by the following Cayley table:

$\circ$	$0$	$a$	$b$	$c$
$0$	$\{0\}$	$\{0\}$	$\{0\}$	$\{0\}$
$a$	$\{a\}$	$\{0\}$	$\{0\}$	$\{0\}$
$b$	$\{b\}$	$\{a\}$	$\{0, a\}$	$\{0, a\}$
$c$	$\{c\}$	$\{a, b, c\}$	$\{a, b, c\}$	$\{0, b, c\}$

Table a3

Then  $(H, \circ, 0)$  is a Smarandache hyper  $K$ -algebra because  $(\Omega = \{0, a, b\}, \circ, 0)$  is a hyper  $BCK$ -algebra.

**Example 3.6.** Let  $H = \{0, a, b\}$  and define an hyper operation “ $\circ$ ” on  $H$  by the following Cayley table:

$\circ$	$0$	$a$	$b$
$0$	$\{0\}$	$\{0\}$	$\{0\}$
$a$	$\{a, b\}$	$\{0, a, b\}$	$\{0, a\}$
$b$	$\{b\}$	$\{a, b\}$	$\{0, a, b\}$

Table a4

Then  $(H, \circ, 0)$  is not a Smarandache hyper  $K$ -algebra since  $(\Omega_1 = \{0, a\}, \circ, 0)$  and  $(\Omega_2 = \{0, b\}, \circ, 0)$  are not hyper  $BCK$ -algebras.

In what follows, let  $H$  and  $\Omega$  denote a Smarandache hyper  $K$ -algebra and a non-trivial hyper  $BCK$ -algebra which is properly contained in  $H$ , respectively.

**Definition 3.7.** A nonempty subset  $I$  of  $H$  is called a *Smarandache hyper  $(\ll, \in)$ -ideal* of  $H$  related to  $\Omega$  (or briefly,  *$\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal* of  $H$ ) if it satisfies:

- (c1)  $0 \in I$ ,
- (c2)  $(\forall x \in \Omega) (\forall y \in I) (x \circ y \ll I \Rightarrow x \in I)$ .

If  $I$  is a Smarandache hyper  $(\ll, \in)$ -ideal of  $H$  related to every hyper  $BCK$ -algebra contained in  $H$ , we simply say that  $I$  is a *Smarandache hyper  $(\ll, \in)$ -ideal* of  $H$ .

**Example 3.8.** Let  $H = \{0, a, b, c\}$  and define a hyper operation “ $\circ$ ” on  $H$  by the following Cayley table:

$\circ$	$0$	$a$	$b$	$c$
$0$	$\{0\}$	$\{0\}$	$\{0\}$	$\{0\}$
$a$	$\{a\}$	$\{0\}$	$\{0\}$	$\{0\}$
$b$	$\{b\}$	$\{b\}$	$\{0\}$	$\{b\}$
$c$	$\{c\}$	$\{a, b\}$	$\{0, a\}$	$\{0, b\}$

Table 3

Then  $(H, \circ, 0)$  is a Smarandache hyper  $K$ -algebra because  $(\Omega = \{0, b\}, \circ, 0)$  is a hyper  $BCK$ -algebra. Moreover, a subset  $\{0, c\}$  of  $H$  is an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal of  $H$ .

**Definition 3.9.** A nonempty subset  $I$  of  $H$  is called a *Smarandache hyper  $(<, \in)$ -ideal* of  $H$  related to  $\Omega$  (or briefly,  *$\Omega$ -Smarandache hyper  $(<, \in)$ -ideal* of  $H$ ) if it satisfies:

- (c1)  $0 \in I$ ,
- (c2\*)  $(\forall x \in \Omega) (\forall y \in I) (x \circ y < I \Rightarrow x \in I)$ .

If  $I$  is a Smarandache hyper  $(<, \in)$ -ideal of  $H$  related to every hyper  $BCK$ -algebra contained in  $H$ , we simply say that  $I$  is a *Smarandache hyper  $(<, \in)$ -ideal* of  $H$ .

**Example 3.10.** Consider a Smarandache hyper  $K$ -algebra  $H = \{0, a, b, c\}$  described in Example 3.8. Then a subset  $\{0, a\}$  of  $H$  is an  $\Omega$ -Smarandache hyper  $(\langle, \in)$ -ideal of  $H$ , where  $\Omega = \{0, a, b\}$ . Moreover, a subset  $\{0, c\}$  of  $H$  is an  $\Omega$ -Smarandache hyper  $(\langle, \in)$ -ideal of  $H$ , where  $\Omega = \{0, b\}$ .

**Theorem 3.11.** Every  $\Omega$ -Smarandache hyper  $(\langle, \in)$ -ideal is a  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal.

*Proof.* Straightforward.  $\square$

**Proposition 3.12.** Let  $A$  be a subset of  $H$  and let  $I$  be an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal of  $H$  which is contained in  $\Omega$ . If  $A \subseteq \Omega$  and  $A \ll I$ , then  $A$  is contained in  $I$ .

*Proof.* Assume that  $A \subseteq \Omega$  and  $A \ll I$ . Let  $x \in A$ . Then  $x \circ 0 = \{x\} \ll I$ , and so  $x \in I$  by (c2). Therefore  $A$  is contained in  $I$ .  $\square$

**Proposition 3.13.** Let  $H$  be a Smarandache hyper  $K$ -algebra with  $\Omega$  as a hyper  $BCK$ -algebra. Then

$$(\forall x, y, z, u \in \Omega) (((x \circ z) \circ (y \circ z)) \circ u \ll (x \circ y) \circ u).$$

*Proof.* Let  $x, y, z, u \in \Omega$ . Since  $\Omega$  is a hyper  $BCK$ -algebra, we have

$$\begin{aligned} ((x \circ z) \circ (y \circ z)) \circ u &= ((x \circ u) \circ z) \circ (y \circ z) \\ &= \cup \{(w \circ z) \circ (y \circ z) \mid w \in x \circ u\} \\ &\ll \cup \{w \circ y \mid w \in x \circ u\} = (x \circ u) \circ y = (x \circ y) \circ u, \end{aligned}$$

which is the desired result.  $\square$

**Definition 3.14.** A nonempty subset  $I$  of  $H$  is called a Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$  related to  $\Omega$  (or briefly,  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$ ) if it satisfies:

- (c1)  $0 \in I$ ,
- (cw)  $(\forall x \in \Omega) (\forall y \in I) (x \circ y \subseteq I \Rightarrow x \in I)$ .

If  $I$  is a Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$  related to every hyper  $BCK$ -algebra contained in  $H$ , we simply say that  $I$  is a Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$ .

**Example 3.15.** Let  $H = \{0, a, b, c\}$  and define an hyper operation “ $\circ$ ” on  $H$  by the following Cayley table:

$\circ$	0	$a$	$b$	$c$
0	$\{0\}$	$\{0, a\}$	$\{0\}$	$\{0\}$
$a$	$\{a\}$	$\{0\}$	$\{a\}$	$\{a\}$
$b$	$\{b\}$	$\{b\}$	$\{0, b\}$	$\{0, b\}$
$c$	$\{c\}$	$\{c\}$	$\{b, c\}$	$\{0, b, c\}$

Table a5

Then  $(H, \circ, 0)$  is a Smarandache hyper  $K$ -algebra because  $(\Omega = \{0, b, c\}, \circ, 0)$  is a hyper  $BCK$ -algebra. Moreover, a subset  $\{0, b\}$  of  $H$  is an  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$ .

**Theorem 3.16.** Every Smarandache hyper  $(\ll, \in)$ -ideal is a Smarandache hyper  $(\subseteq, \in)$ -ideal.

*Proof.* Note that  $A \subseteq B \Rightarrow A \ll B$  for every subsets  $A$  and  $B$  of  $H$ . Thus the result is obvious.  $\square$

The converse of Theorem 3.16 is not true in general as seen in the following example.

**Example 3.17.** Consider Example 3.15. Then  $\{0, c\}$  is an  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$  which is not an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal of  $H$ , where  $\Omega = \{0, b, c\}$ .

**Theorem 3.18.** Let  $\Omega_1$  and  $\Omega_2$  be hyper BCK-algebras which are properly contained in  $H$  such that  $\Omega_1 \subset \Omega_2$ . Then every  $\Omega_2$ -Smarandache hyper  $(\ll, \in)$ -ideal is an  $\Omega_1$ -Smarandache hyper  $(\ll, \in)$ -ideal.

*Proof.* Straightforward. □

**Theorem 3.19.** Let  $\Omega_1$  and  $\Omega_2$  be hyper BCK-algebras which are properly contained in  $H$  such that  $\Omega_1 \subset \Omega_2$ . Then every  $\Omega_2$ -Smarandache hyper  $(<, \in)$ -ideal is an  $\Omega_1$ -Smarandache hyper  $(<, \in)$ -ideal.

*Proof.* Straightforward. □

**Theorem 3.20.** Let  $\Omega_1$  and  $\Omega_2$  be hyper BCK-algebras which are properly contained in  $H$  such that  $\Omega_1 \subset \Omega_2$ . Then every  $\Omega_2$ -Smarandache hyper  $(\subseteq, \in)$ -ideal is an  $\Omega_1$ -Smarandache hyper  $(\subseteq, \in)$ -ideal.

*Proof.* Straightforward. □

**Definition 3.21.** A nonempty subset  $I$  of  $H$  is called a Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$  related to  $\Omega$  (or briefly,  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$ ) if it satisfies:

- (c1)  $0 \in I$ ,
- (cs)  $(\forall x, y, z \in \Omega) ((x \circ y) \circ z \ll I, y \circ z \subseteq I \Rightarrow x \circ z \subseteq I)$ .

**Theorem 3.22.** Every  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal is an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal.

*Proof.* Let  $I$  be an  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$  and assume that  $x \circ y \ll I$  for all  $x \in \Omega$  and  $y \in I$ . Taking  $z = 0$  in (cs), we have  $(x \circ y) \circ 0 = x \circ y \ll I$  and  $y \circ 0 = \{y\} \subseteq I$ . Using (cs), we conclude that  $\{x\} = x \circ 0 \subseteq I$ . Hence  $I$  is an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal of  $H$ . □

The following example shows that the converse of Theorem 3.22 is not true in general.

**Example 3.23.** Let  $H = \{0, a, b, c\}$  and define an hyper operation “ $\circ$ ” on  $H$  by the following Cayley table:

$\circ$	0	a	b	c
0	{0}	{0, a}	{0}	{0}
a	{a}	{0, a}	{a}	{a}
b	{b}	{b}	{0}	{0}
c	{c}	{c}	{b}	{0, b}

Table a6

Then  $(H, \circ, 0)$  is a Smarandache hyper  $K$ -algebra because  $(\Omega = \{0, b, c\}, \circ, 0)$  is a hyper BCK-algebra. Moreover, a subset  $\{0\}$  of  $H$  is an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal of  $H$  which is not an  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$ .

**Theorem 3.24.** If  $I$  is an  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$ , then the set

$$\Omega(I_a) := \{x \in \Omega \mid x \circ a \subseteq I\}$$

is an  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$  for all  $a \in H$ .

*Proof.* Obviously  $0 \in \Omega(I_a)$ . Assume that  $x \circ y \subseteq \Omega(I_a)$  for all  $x \in \Omega$  and  $y \in \Omega(I_a)$ . Then  $(x \circ y) \circ a \subseteq I$ , and hence  $(x \circ y) \circ a \ll I$ , and  $y \circ a \subseteq I$ . It follows from (cs) that  $x \circ a \subseteq I$ , that is,  $x \in \Omega(I_a)$ . Therefore  $\Omega(I_a)$  is an  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$ . □

**Theorem 3.25.** *Let  $I$  be an  $\Omega$ -Smarandache hyper  $(\ll, \in)$ -ideal of  $H$  which is contained in  $\Omega$  such that the set*

$$\Omega(I_a) := \{x \in \Omega \mid x \circ a \subseteq I\}, a \in H$$

*is an  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal of  $H$ . Then  $I$  is an  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$ .*

*Proof.* Assume that  $(x \circ y) \circ z \ll I$  and  $y \circ z \subseteq I$  for all  $x, y, z \in \Omega$ . From  $y \circ z \subseteq I$ , we get  $y \in \Omega(I_z)$ . Since  $(x \circ y) \circ z \subseteq \Omega$ , it follows from Proposition 3.12 that  $(x \circ y) \circ z \subseteq I$  so that  $w \circ z \subseteq I$ , that is,  $w \in \Omega(I_z)$  for each  $w \in x \circ y$ . Hence  $x \circ y \subseteq \Omega(I_z)$ . Since  $\Omega(I_z)$  is an  $\Omega$ -Smarandache hyper  $(\subseteq, \in)$ -ideal, we have  $x \in \Omega(I_z)$ , and so  $x \circ z \subseteq I$ . Therefore  $I$  is an  $\Omega$ -Smarandache hyper  $(\ll, \subseteq, \subseteq)$ -ideal of  $H$ .  $\square$

#### 4. ACKNOWLEDGEMENTS

The author Y. B. Jun was supported by Korea Research Foundation Grant (KRF-2003-005-C00013).

#### REFERENCES

- [1] R. A. Borzoei, A. Hasankhani, M. M. Zahedi and Y. B. Jun, *On hyper  $K$ -algebras*, Math. Japonica **52(1)** (2000), 113–121.
- [2] Y. B. Jun, *Smarandache BCI-algebras*, Sci. Math. Japon. (submitted).
- [3] J. Meng and Y. B. Jun, *BCK-algebras*, Kyungmoonsa Co. Seoul, Korea (1994).
- [4] R. Padilla, *Smarandache algebraic structures*, Bull. Pure Appl. Sci., Delhi, **17E** (1998), no. 1, 119–121; <http://www.gallup.unm.edu/smarandache/alg-s-tx.txt>.
- [5] W. B. Vasantha Kandasamy, *Smarandache groupoids*, <http://www.gallup.unm.edu/smarandache/Groupoids.pdf>.
- [6] M. M. Zahedi, R. A. Borzoei, Y. B. Jun and A. Hasankhani, *Some results on hyper  $K$ -algebras*, Scientiae Mathematicae **3(1)** (2000), 53–59.

Y. B. JUN, DEPARTMENT OF MATHEMATICS EDUCATION, GYEONGSANG NATIONAL UNIVERSITY, JINJU 660-701, KOREA

*E-mail address:* ybjun@gsnu.ac.kr

E. H. ROH, DEPARTMENT OF MATHEMATICS EDUCATION, CHINJU NATIONAL UNIVERSITY OF EDUCATION, JINJU 660-756, KOREA

*E-mail address:* ehroh@cue.ac.kr